

Horn Torus Model for the Riemann Sphere From the Viewpoint of Division by Zero (memo)

Wolfgang W. Däumler, Perouse, Germany,

Vyacheslav Puha

Kola Science Centre of the Russian Academy of Sciences

14, Fersman str., Apatity, Murmansk reg., RUSSIA 184209 and

Saburou Saitoh

Institute of Reproducing Kernels

Kawauchi-cho, 5-1648-16, Kiryu 376-0041, JAPAN

September 16, 2018

Abstract: In this paper, we will introduce a beautiful horn torus model for the Riemann sphere in complex analysis from the viewpoint of the division by zero.

Key Words: Division by zero, division by zero calculus, singular point, $0/0 = 1/0 = z/0 = 0$, infinity, discontinuous, point at infinity, stereographic projection, Riemann sphere, horn torus, Laurent expansion.

1 Division by zero calculus and introduction

The division by zero with mysterious and long history was indeed trivial and clear as in the followings:

By the concept of the Moore-Penrose generalized solution of the fundamental equation $ax = b$, the division by zero was trivial and clear all as

$a/0 = 0$ in the **generalized fraction** that is defined by the generalized solution of the equation $ax = b$.

Division by zero is trivial and clear from the concept of repeated subtraction - H. Michiwaki.

Recall the uniqueness theorem by S. Takahasi on the division by zero.

The simple field structure containing division by zero was established by M. Yamada.

Many applications of the division by zero to Wasan geometry were given by H. Okumura.

The division by zero opens a new world since Aristotelēs-Euclid. See the references for recent related results.

As the number system containing the division by zero, the Yamada field structure is complete.

However, for applications of the division by zero to **functions**, we need the concept of the division by zero calculus for the sake of uniquely determinations of the results and for other reasons.

For example, for the typical linear mapping

$$W = \frac{z - i}{z + i}, \quad (1.1)$$

it gives a conformal mapping on $\{\mathbf{C} \setminus \{-i\}\}$ onto $\{\mathbf{C} \setminus \{1\}\}$ in one to one and from

$$W = 1 + \frac{-2i}{z - (-i)}, \quad (1.2)$$

we see that $-i$ corresponds to 1 and so the function maps the whole $\{\mathbf{C}\}$ onto $\{\mathbf{C}\}$ in one to one.

Meanwhile, note that for

$$W = (z - i) \cdot \frac{1}{z + i}, \quad (1.3)$$

we should not enter $z = -i$ in the way

$$[(z - i)]_{z=-i} \cdot \left[\frac{1}{z + i} \right]_{z=-i} = (-2i) \cdot 0 = 0. \quad (1.4)$$

However, in many cases, the above two results will have practical meanings and so, we will need to consider many ways for the application of the

division by zero and we will need to check the results obtained, in some practical viewpoints. We referred to this delicate problem with many examples.

Therefore, we will introduce the division by zero calculus. For any Laurent expansion around $z = a$,

$$f(z) = \sum_{n=-\infty}^{-1} C_n(z-a)^n + C_0 + \sum_{n=1}^{\infty} C_n(z-a)^n, \quad (1.5)$$

we obtain the identity, by the division by zero

$$f(a) = C_0. \quad (1.6)$$

Note that here, there is no problem on any convergence of the expansion (1.5) at the point $z = a$, because all the terms $(z-a)^n$ are zero at $z = a$ for $n \neq 0$.

For the correspondence (1.6) for the function $f(z)$, we will call it **the division by zero calculus**. By considering the formal derivatives in (1.5), we can define any order derivatives of the function f at the singular point a ; that is,

$$f^{(n)}(a) = n!C_n.$$

Apart from the motivation, we define the division by zero calculus by (1.6). With this assumption, we can obtain many new results and new ideas. However, for this assumption we have to check the results obtained whether they are reasonable or not. By this idea, we can avoid any logical problems. – In this point, the division by zero calculus may be considered as a fundamental assumption like an axiom.

For the fundamental function $W = 1/z$ we did not consider any value at the origin $z = 0$, because we did not consider the division by zero $1/0$ in a good way. Many and many people consider its value by the limiting like $+\infty$ and $-\infty$ or the point at infinity as ∞ . However, their basic idea comes from **continuity** with the common sense or based on the basic idea of Aristotle. – For the related Greece philosophy, see [19, 20, 21]. However, as the division by zero we will consider its value of the function $W = 1/z$ as zero at $z = 0$. We will see that this new definition is valid widely in mathematics and mathematical sciences, see ([6, 7]) for example. Therefore, the division by zero will give great impacts to calculus, Euclidian geometry, analytic geometry, complex analysis and the theory of differential equations

in an undergraduate level and furthermore to our basic ideas for the space and universe.

For the extended complex plane, we consider its stereographic projection mapping as the Riemann sphere and the point at infinity is realized as the north pole in the Alexandroff's one point compactification. The Riemann sphere model gives a beautiful and complete realization of the extended complex plane through the stereographic projection mapping and the mapping has beautiful properties like isogonal (equiangular) and circle to circle correspondence (circle transformation). Therefore, the Riemann sphere is a very classical concept [1].

Now, with the division by zero we have to admit the strong discontinuity at the point at infinity, because the point at infinity is represented by zero. In [7], a formal contradiction for the classical result $1/0 = \infty$ was given and the strong discontinuity was shown in many and many examples. See the papers in the references.

On this situation, the second author discovered the mapping of the extended complex plane to a beautiful horn torus at (2018.6.4.7:22) and its inverse at (2018.6.18.22:18).

Incidentally, independently of the division by zero, Wolfgang W. Daeumler has various special great ideas on horn torus as we see from his site:

Horn Torus & Physics (<https://www.horntorus.com/>) 'Geometry Of Everything', intellectual game to reveal engrams of dimensional thinking and proposal for a different approach to physical questions ...

Indeed, Wolfgang Daeumler was presumably the first (1996) who came to the idea of the possibility of a mapping of extended complex plane onto the horn torus. He expressed this idea on his private website (<http://www.dorntorus.de>). He was also, apparently, the first who to point out that zero and infinity are represented by one and the same point on the horn torus model of extended complex plane.

In this paper we just introduce a new horn torus model for the classical Riemann sphere from the viewpoint of the division by zero. This model seems to be important for us.

2 Horn torus model

We will consider the three circles stated by

$$\begin{aligned}\xi^2 + \left(\zeta - \frac{1}{2}\right)^2 &= \left(\frac{1}{2}\right)^2, \\ \left(\xi - \frac{1}{4}\right)^2 + \left(\zeta - \frac{1}{2}\right)^2 &= \left(\frac{1}{4}\right)^2,\end{aligned}\tag{2.1}$$

and

$$\left(\xi + \frac{1}{4}\right)^2 + \left(\zeta - \frac{1}{2}\right)^2 = \left(\frac{1}{4}\right)^2.$$

By rotation on the space (ξ, η, ζ) on the (x, y) plane as in $\xi = x, \eta = y$ around ζ axis, we will consider the sphere with $1/2$ radius as the Riemann sphere and the horn torus made in the sphere.

The stereographic projection mapping from (x, y) plane to the Riemann sphere is given by

$$\begin{aligned}\xi &= \frac{x}{x^2 + y^2 + 1}, \\ \eta &= \frac{y}{x^2 + y^2 + 1},\end{aligned}$$

and

$$\zeta = \frac{x^2 + y^2}{x^2 + y^2 + 1}.$$

Of course,

$$\xi^2 + \eta^2 = \zeta(1 - \zeta).$$

and

$$x = \frac{\xi}{1 - \zeta}, y = \frac{\eta}{1 - \zeta}.$$

The mapping from (x, y) plane to the horn torus is given by

$$\begin{aligned}\xi &= \frac{2x\sqrt{x^2 + y^2}}{(x^2 + y^2 + 1)^2}, \\ \eta &= \frac{2y\sqrt{x^2 + y^2}}{(x^2 + y^2 + 1)^2},\end{aligned}$$

and

$$\zeta = \frac{(x^2 + y^2 - 1)\sqrt{x^2 + y^2}}{(x^2 + y^2 + 1)^2} + \frac{1}{2}.$$

The inversion is given by

$$x = \xi \left(\xi^2 + \eta^2 + \left(\zeta - \frac{1}{2} \right)^2 - \zeta + \frac{1}{2} \right)^{(-1/2)} \quad (2.2)$$

and

$$y = \eta \left(\xi^2 + \eta^2 + \left(\zeta - \frac{1}{2} \right)^2 - \zeta + \frac{1}{2} \right)^{(-1/2)}. \quad (2.3)$$

In these formulas, we can see the division by zero

$$\frac{1}{0} = 0,$$

naturally.

3 Properties of horn torus model

At first, the model shows the strong symmetry of the domains $\{|z| < 1\}$ and $\{|z| > 1\}$ and they correspond to the lower part and the upper part of the horn torus, respectively. The unit circle $\{|z| = 1\}$ corresponds to the circle

$$\xi^2 + \eta^2 = \left(\frac{1}{2} \right)^2, \quad \zeta = \frac{1}{2}$$

in one to one way. Of course, the origin and the point at infinity are the same point and correspond to $(0, 0, 1/2)$. Furthermore, the inversion relation

$$z \longleftrightarrow \frac{1}{\bar{z}}$$

with respect to the unit circle $\{|z| = 1\}$ corresponds to the relation

$$(\xi, \eta, \zeta) \longleftrightarrow (\xi, \eta, 1 - \zeta)$$

and similarly,

$$z \longleftrightarrow -z$$

corresponds to the relation

$$(\xi, \eta, \zeta) \longleftrightarrow (-\xi, -\eta, \zeta)$$

and

$$z \longleftrightarrow -\frac{1}{\bar{z}}$$

corresponds to the relation

$$(\xi, \eta, \zeta) \longleftrightarrow (-\xi, -\eta, 1 - \zeta)$$

(H.G.W. Begehr: 2018.6.18.19:20).

We can see directly the important negative properties that the mapping is not isogonal (equiangular) and infinitely small circles do not correspond to infinitely small circles, as in analytic functions.

Horn torus, in contrast to the Riemann sphere, does not satisfy the definition of simply connected space because a closed nonzero path passing through the point $(0, 0, 1/2)$ can not be continuously shrunk to the point. In particular, note that a curve can pass the point $(0, 0, 1/2)$ on the horn torus.

We note that only zero and numbers a of the form $|a| = 1$ have the property : $|a|^b = |a|, b \neq 0$. Here, note that we can also consider $0^b = 0$ ([4]). The symmetry of the horn torus model agrees perfectly with this fact. Only zero and numbers a of the form $|a| = 1$ correspond to points on the plane described by equation $\zeta - 1/2 = 0$. Only zero and numbers a of the form $|a| = 1$ correspond to points whose tangent lines to the surface of the horn torus are parallel to the axis ζ .

4 Conformal mapping from the plane to the horn torus with a modified mapping

W. W. Däumle discovered a suprising conformal mapping from the extended complex plane to the horn torus model (2018.8.18.09):

<https://www.horntorus.com/manifolds/conformal.html>

and

<https://www.horntorus.com/manifolds/solution.html>

Our situation is invariant by rotation around ζ axis, and so we shall consider the problem on the ξ, ζ plane.

Let $N(0, 0, 1)$ be the north pole. Let $P'(\xi, \eta, \zeta)$ denote a point on the horn torus and let $z = x + iy$ be the common point with the line NP' and $\zeta = 0$ plane ($: z = x + iy$); that is P' is the stereographic projection map of the point $z = x + iy$ onto the unit sphere.

Let $M(1/4, 0, 1/2)$ be the center of the circle (2.1). Let P'' be the common point of the line $SP'(S = S(0, 0, 1/2))$ and the circle (2.1).

Let Q' be $(0, 0, \zeta)$ that is the line $Q'P'$ is parallel to the x axis. Let Q'' and M'' be the common points with the ζ axis and $\xi = 1/4$ with the parallel line to the x axis through the point P'' , respectively.

Further, we set $\alpha = \angle OSP' = \angle P''IS = (1/2)\angle P''MS$ ($I := I(1/2, 0, 1/2)$). We set P for the point on the horn torus such that $\phi = \angle SMP$ and Q be the point on the ζ axis such that the line QP is parallel to the x axis.

Then, we have:

$$\begin{aligned}\overline{P'Q'} &= \frac{1}{2} \sin \alpha, \\ \overline{P''M''} &= \frac{1}{4} |\cos(2\alpha)|, \\ \overline{P''Q''} &= \frac{1}{4} (1 - \cos(2\alpha)),\end{aligned}$$

the length of latitude through P' is

$$2\pi \overline{P'Q'} = \pi \sin \alpha,$$

and the length of latitude through P''

$$2\pi \overline{P''Q''} = \frac{\pi}{2} (1 - \cos(2\alpha)) = \pi \sin^2 \alpha.$$

Similarly, we have

$$2\pi \overline{QP} = \frac{\pi}{2} (1 - \cos \phi).$$

We observe a small circle around P' on the sphere with radius $d\alpha$, the radius perpendicular to screen in P' with same length is $d\omega = d\alpha / \sin \alpha$ (ω is longitude) the projected small circle around P on horn torus has the same radius $d\omega$ (as angle, not length) and for corresponding perpendicular radius ϕ we respectively have: $d\omega = 2d\phi / (1 - \cos \phi)$, ω and $d\omega$, as angles, are identical on both figures, therefore:

$$\frac{d\alpha}{\sin \alpha} = \frac{2d\phi}{1 - \cos \phi}$$

that ensures the conformal correspondence between P' and P .

We solve this differential equation as

$$\log \left| \tan \frac{\alpha}{2} \right| = -\cot \frac{\phi}{2} + C,$$

using the correspondence

$$\alpha = \pi/2 \leftrightarrow \phi = \pi$$

or

$$\alpha = 0 \leftrightarrow \phi = 0,$$

or

$$\alpha = \pi \leftrightarrow 2\phi = 0,$$

we have $C = 0$. Note that $\log 0 = 0$ ([4]). By solving for ϕ we have the result

$$\phi = 2 \cot^{-1}(-\log \left| \tan(\alpha/2) \right|) \quad (4.1)$$

or

$$\alpha = 2 \tan^{-1}(e^{(-\cot(\phi/2))}). \quad (4.2)$$

Next, note that

$$\tan \frac{\alpha}{2} = |z|$$

and

$$\alpha = 2 \tan^{-1} |z|. \quad (4.3)$$

We thus have

$$\phi = 2 \cot^{-1}(-\log |z|) \quad (4.4)$$

and the inverse is

$$|z| = e^{-\cot(\phi/2)}. \quad (4.5)$$

We thus obtain the complicated conformal mapping for the z plane to the horn torus by (4.4) and (4.2). The converse conformal mapping for the horn torus to the complex z plane is given by (4.1) and (4.5).

We can represent the direct Däumler mapping from the z plane onto the horn torus as follows (V. Puha: 2018.8.28.22:31):

$$\xi = \frac{x \cdot (1/2)(\sin(\phi/2))^2}{\sqrt{x^2 + y^2}}, \quad (4.6)$$

$$\eta = \frac{y \cdot (1/2)(\sin(\phi/2))^2}{\sqrt{x^2 + y^2}}, \quad (4.7)$$

and

$$\zeta = -\frac{1}{4} \sin \phi + \frac{1}{2}. \quad (4.8)$$

Indeed, at first, we have

$$SP := L = 2 \cdot \frac{1}{4} \sin \frac{\phi}{2} = \frac{1}{2} \sin \frac{\phi}{2},$$

$$\sqrt{\xi^2 + \eta^2} = L \cos \left(\frac{\pi}{2} - \frac{\phi}{2} \right) = L \sin \frac{\phi}{2},$$

and

$$\sqrt{\xi^2 + \eta^2} = \frac{1}{2} \sin^2 \frac{\phi}{2}.$$

From the simple relations

$$\xi = \frac{x\sqrt{\xi^2 + \eta^2}}{\sqrt{x^2 + y^2}}, \eta = \frac{y\sqrt{\xi^2 + \eta^2}}{\sqrt{x^2 + y^2}},$$

and

$$\zeta = -T \sin \left(\frac{\pi}{2} - \frac{\phi}{2} \right) + \frac{1}{2}$$

we have the desired representations.

5 Acknowledgements

The third author wishes to express their deep thanks Professors H.G.W. Begehr and Tao Qian for their exciting communications.

References

- [1] L. V. Ahlfors, Complex Analysis, McGraw-Hill Book Company, 1966.
- [2] M. Kuroda, H. Michiwaki, S. Saitoh, and M. Yamane, New meanings of the division by zero and interpretations on $100/0 = 0$ and on $0/0 = 0$, Int. J. Appl. Math. **27** (2014), no 2, pp. 191-198, DOI: 10.12732/ijam.v27i2.9.

- [3] T. Matsuura and S. Saitoh, Matrices and division by zero $z/0 = 0$, *Advances in Linear Algebra & Matrix Theory*, **6**(2016), 51-58 Published Online June 2016 in SciRes. <http://www.scirp.org/journal/alamt> <http://dx.doi.org/10.4236/alamt.2016.62007>.
- [4] T. Matsuura, H. Michiwaki and S. Saitoh, $\log 0 = \log \infty = 0$ and applications. *Differential and Difference Equations with Applications. Springer Proceedings in Mathematics & Statistics*. **230** (2018), 293-305.
- [5] H. Michiwaki, S. Saitoh and M. Yamada, Reality of the division by zero $z/0 = 0$. *IJAPM International J. of Applied Physics and Math.* **6**(2015), 1-8. <http://www.ijapm.org/show-63-504-1.html>
- [6] H. Michiwaki, H. Okumura and S. Saitoh, Division by Zero $z/0 = 0$ in Euclidean Spaces, *International Journal of Mathematics and Computation*, **28**(2017); Issue 1, 1-16.
- [7] H. Okumura, S. Saitoh and T. Matsuura, Relations of 0 and ∞ , *Journal of Technology and Social Science (JTSS)*, **1**(2017), 70-77.
- [8] H. Okumura and S. Saitoh, The Descartes circles theorem and division by zero calculus. <https://arxiv.org/abs/1711.04961> (2017.11.14).
- [9] H. Okumura, Wasan geometry with the division by 0. <https://arxiv.org/abs/1711.06947> *International Journal of Geometry*.
- [10] H. Okumura and S. Saitoh, Harmonic Mean and Division by Zero, Dedicated to Professor Josip Pećarić on the occasion of his 70th birthday, *Forum Geometricorum*, **18** (2018), 155—159.
- [11] H. Okumura and S. Saitoh, Remarks for The Twin Circles of Archimedes in a Skewed Arbelos by H. Okumura and M. Watanabe, *Forum Geometricorum*, **18**(2018), 97-100.
- [12] H. Okumura and S. Saitoh, Applications of the division by zero calculus to Wasan geometry. *GLOBAL JOURNAL OF ADVANCED RESEARCH ON CLASSICAL AND MODERN GEOMETRIES*” (GJARC-MG)(in press).

- [13] S. Pinelas and S. Saitoh, Division by zero calculus and differential equations. *Differential and Difference Equations with Applications. Springer Proceedings in Mathematics & Statistics.* **230** (2018), 399-418.
- [14] S. Saitoh, Generalized inversions of Hadamard and tensor products for matrices, *Advances in Linear Algebra & Matrix Theory.* **4** (2014), no. 2, 87–95. <http://www.scirp.org/journal/ALAMT/>
- [15] S. Saitoh, A reproducing kernel theory with some general applications, Qian,T./Rodino,L.(eds.): *Mathematical Analysis, Probability and Applications - Plenary Lectures: Isaac 2015, Macau, China, Springer Proceedings in Mathematics and Statistics,* **177**(2016), 151-182.
- [16] S. Saitoh, Mysterious Properties of the Point at Infinity, arXiv:1712.09467 [math.GM](2017.12.17).
- [17] S. Saitoh, Division by zero calculus (189 pages): <http://okmr.yamatoblog.net/>
- [18] S.-E. Takahasi, M. Tsukada and Y. Kobayashi, Classification of continuous fractional binary operations on the real and complex fields, *Tokyo Journal of Mathematics,* **38**(2015), no. 2, 369-380.
- [19] <https://philosophy.kent.edu/OPA2/sites/default/files/012001.pdf>
- [20] http://publish.uwo.ca/~jbell/The_20Continuous.pdf
- [21] <http://www.mathpages.com/home/kmath526/kmath526.htm>

**E-mail: artmetic@gmx.de, puhanot@gmail.com,
saburou.saitoh@gmail.com**